

Fibromyalgia: clinical and occupational aspects

MILTON HELFENSTEIN JUNIOR¹, MARCO AURÉLIO GOLDENFUM², CÉSAR AUGUSTO FÁVARO SIENA³

¹ PhD in Rheumatology; Assistant Doctor, Universidade Federal de São Paulo (Unifesp), São Paulo, SP, Brazil

² MSc Student in University Teaching, National Technological University of Buenos Aires, Argentina; Coordinator, Postgraduate Program in Occupational Medicine and Medical Inspections, Escola Superior de Gestão e Ciências da Saúde de Porto Alegre, Porto Alegre, RS, Brazil

³ Federal Court Medical Expert of Ribeirão Preto, Ribeirão Preto, SP, Brazil

SUMMARY

Fibromyalgia (FM) is a clinical syndrome commonly observed in daily medical practice and its etiopathogenesis is still unclear. As it is characterized by chronic musculoskeletal pain associated with several symptoms, FM may be confused with several other rheumatic and nonrheumatic diseases when they course with pictures of diffuse pain and chronic fatigue. FM treatment should be multidisciplinary, individualized, count on active participation of the patient, and based on combined pharmacological and non-pharmacological modalities. It is found both in work and non-work settings, and there is no scientific evidence in the literature showing that FM might be caused by occupation. FM seldom leads to incapacity to work. In cases where pain or fatigue do not respond to appropriate treatment, reaching significant levels, a short period away from work can be considered. As FM is a relevant subject, this review article was based on exploratory, qualitative and bibliographic investigation, aiming to study the main clinical and occupational aspects of FM, emphasizing the theoretical-conceptual background and the experience of specialists.

Keywords: Fibromyalgia; review; rheumatology; occupational medicine; referred pain.

©2012 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de [CC BY-NC-ND](#)

Study conducted at the Discipline
of Rheumatology, Escola Paulista
de Medicina, Universidade
Federal de São Paulo
São Paulo, SP, Brazil

Submitted on: 10/20/2011
Approved on: 02/10/2012

Correspondence to:
Milton Helfenstein Junior
Rua João de Lacerda Soares, 90
Brooklin
São Paulo – SP, Brazil
CEP: 04707-010
m.helfen@terra.com.br

Conflicts of interest: None.

INTRODUCTION

Fibromyalgia (FM) can be described as a painful clinical syndrome associated with other symptoms, and it is also termed fibromyalgia syndrome. In this review, the term fibromyalgia, used in the International Classification of Diseases (ICD), has been adopted.

Although it has been recognized for a long time, FM has been seriously studied for the last four decades. FM was not considered a well-defined clinical entity until the 1970's, when the first reports on sleep disturbances were published. The concept of FM was introduced in 1977, when anatomical sites with excessive pain sensitivity were described; they were termed tender points. Sleep disturbances, including those experimentally induced, were also documented, possibly reproducing pain and muscular sensitivity symptoms seen in this chronic pain syndrome¹.

The term fibromyalgia is derived from the Latin word *fibro* (fibrous tissue, present in ligaments, tendons and fasciae) and from the Greek words *mio* (muscular tissue), *algos* (pain), and *ia* (condition). It was first proposed by Yunun et al. in 1981, aiming to replace the term fibrositis², hitherto used to name a particular type of rheumatism characterized by the presence of tender indurated muscular points, which were understood as having no tissue inflammation³.

In the 1980's, several diagnostic criteria based on systemic disease exclusion, presence of certain symptoms, and tender points on physical examination were suggested.

In 1990, a committee from the American College of Rheumatology (ACR) defined the FM classification criteria as the presence of a clinical history of widespread pain, affecting the axial and peripheral skeleton, above and below the waist, and with a duration > 3 months; physical examination pain elicited by application of a 4 kg/cm² force to at least 11 of the following 18 *tender points* (9 pairs): 1) suboccipital muscle insertions; 2) ligaments of intertransverse spaces C5-C7; 3) trapezius upper border; 4) supraspinatus muscle origins; 5) pectoral muscle junction with the second rib costocondral junction; 6) 2 cm distal to the elbow lateral epicondyle; 7) upper outer quadrant of buttocks, below the iliac spine; 8) muscle insertions at the femoral trochanter; 9) fat pad, slightly above the knee midline⁴.

The current diagnostic criteria do not include tender points; however, they consider symptoms not related to the locomotor system. In addition to musculoskeletal pain, they assess the syndrome's severity and are useful to establish the diagnosis⁵.

Considering the subject's relevance, this review aims to discuss the main clinical and occupational aspects of FM.

METHODS

This review article is derived from a qualitative, exploratory and bibliographic investigation performed in the electronic libraries SciELO and PubMed, emphasizing the theoretical-conceptual background and the experience of specialists. The keywords "fibromyalgia", "review", "rheumatology", "occupational medicine", and "referred pain" were searched. Moreover, a manual search of references from selected articles was performed so that papers not mentioned in the databases could be found. The references contributing to meet the search target, i.e., review of the main clinical and occupational aspects of FM, were selected from the material retrieved. The investigation was closed only when signs of search theme theoretical saturation were found.

EPIDEMIOLOGY

FM occurs at any age and is much more often diagnosed in females. A study conducted by the American College of Rheumatology found an FM prevalence of 3.4% for females and 0.5% for males, with an estimated prevalence of 2% for both genders⁶.

The first studies dedicated to FM prevalence were reported in the 1980's, but the diagnostic criteria were different, albeit similar to the current criteria. The results ranged from 2.1% in private practices, 5.7% in outpatient facilities, 5% to 8% for inpatients, and as high as 14% to 20% in rheumatology outpatient settings⁷.

A Brazilian study determined a prevalence of 2.5% in the population, with the majority of cases identified in females, of whom 40.8% were aged 35 to 44 years⁸. In some European countries, FM rates found in the adult population were as high as 10.5%⁹.

FM is a condition commonly seen in daily clinical practice and one of the main causes for medical visits linked to the musculoskeletal system; it is considered the second main rheumatologic disorder, surpassed only by osteoarthritis¹⁰.

Despite the good prognosis for this painful condition, patients with FM consume huge amounts of financial resources in public or private health care both for treatment and diagnostic investigation.

The magnitude of economic and financial impact of FM on society is evidenced in an American study reported in 2007, finding an annual cost of USD 9,573 per patient, and representing an expenditure three to five times higher than that of the general population¹¹.

Likewise, a survey conducted in the Netherlands identified a mean annual cost of € 7,813 per patient with FM, € 8,553 for chronic low back pain, and € 3,205 for ankylosing spondylitis – the latter amount was calculated before biologics were approved for ankylosing spondylitis treatment¹².

Assuming that FM affects 3% of the population, according to statistical data, and using data from demographical studies, these estimates can be translated into an incremental annual cost of about €12 billion for a population of 80 million, with € 960 million (8%) representing the costs of drug therapy¹³.

ETIOPATHOGENESIS

FM etiology and pathophysiology are still nuclear. The current hypotheses focus on the central mechanisms of pain modulation and amplification in the genesis of FM. A pathophysiological model integrating many of the ideas found in literature has been accepted; it suggests that the primary disorder in FM would be a change in some central mechanism of pain control, possibly resulting from a neurotransmitter dysfunction. Such neurohormonal dysfunction would include either an inhibitory neurotransmitter deficiency in spinal or supraspinal levels (serotonin, enkephalin, norepinephrine, and others) or an excitatory neurotransmitter hyperactivity (substance P, glutamate, bradykinin, and other peptides). Both conditions might be present. These dysfunctions could be genetically predetermined and triggered by a nonspecific stress, such as a viral infection, psychological stress, or physical trauma¹⁴.

The hypothalamic-pituitary-adrenal axis and the sympathetic nervous system, comprising the main systems of stress response along with their interactions with neurohormonal dysfunctions, are also implicated in the pathophysiology. Susceptibility to FM development seems to be influenced by environmental, hormonal, and genetic factors^{15,16}, causing changes at the neurohormonal receptors level. An acute stressor could trigger the development of a disordered hypothalamic-pituitary-adrenal axis through unclear mechanisms, possibly involving the sympathetic nervous system and the serotonergic system. Thus, the hypothalamic-pituitary-adrenal axis is believed to play a role in mediating and perpetuating FM symptoms¹⁷.

DIAGNOSIS

FM diagnosis is notably clinical⁵. In addition to diffuse pain in skeletal muscle and the physical examination finding of multiple tender points, most patients with FM also report fatigue, muscle stiffness, pain after exertion, and sleep disturbances. Symptoms of depression, anxiety, memory deficit, inattention, tension headache or migraine, dizziness, vertigo, tingling, symptoms consistent with irritable bowel syndrome or restless feet syndrome, are among several other symptoms not related to the locomotor system¹⁸. Generalized chronic pain, however, is

a cardinal symptom. Myalgia may often show a migratory characteristic responding to biomechanical stress or trauma¹⁹.

Despite having too low diagnostic accuracy to be considered classification criteria, sleep disturbances occur in up to 100% of patients with fibromyalgia and are highly variable. Sleep disturbances are characterized by difficulty in sleep induction, too many awakenings during the night, and a sense that the sleep is not restorative. Sleep disturbances evoke adverse consequences, such as cognitive deficits, morning tiredness, and tendency towards triggering psychiatric disorders^{20,21}.

The psychological profile of patients with fibromyalgia is associated with perfectionism, severe self-criticism, and obsessive search for details²². The high depression prevalence among patients with fibromyalgia is noteworthy²³⁻²⁵.

Depression characteristics, such as fatigue, guilt feelings, low self-esteem, and victimization lead to a symptom flare and impair the patient's coping strategies toward the disease²⁶.

So far, there are no specific laboratory tests to confirm the FM diagnosis. The patients have normal inflammatory activity tests and imaging studies²⁷.

DIFFERENTIAL DIAGNOSIS

FM may be confused with several other diseases when they course with diffuse pain pictures and chronic fatigue.

The patient with hypothyroidism can have a clinical picture mimicking FM. Thus, as these patients are approached, thyroid function assays are recommended. The literature describes FM clinical picture as an initial manifestation of hypothyroidism²⁸. However, there is no evidence that stabilizing thyroid function would dispel the FM picture, and there is no evidence that most patients with FM have a thyroid dysfunction.

Another hormone disorder possibly causing diagnostic confusion with FM is primary hyperparathyroidism. This disorder results from a parathyroid adenoma (monoclonal origin) in 80% to 90% of cases. In nearly all remaining cases, it results from hyperplasia in the four parathyroids (polyclonal origin). In addition to bone (bone pain, pathological fractures, cortical osteopenia, bone cysts, and osteitis fibrosa cystica) and kidney (renal colic, nephrolithiasis, and nephrocalcinosis with kidney failure) manifestations, which are classically part of the disease, unspecific symptoms, such as fatigue, emotional disorder, neuropsychiatric abnormalities and muscular aches may be present, mimicking the clinical picture of FM²⁹.

Vitamin D deficiency with secondary hyperparathyroidism may cause bone and muscular pain sometimes misinterpreted as FM³⁰.

In certain settings, osteomalacia can be part of differential diagnosis, as a number of patients have an onset with weakness, muscular pain, and bone tenderness before the radiologic and metabolic manifestations are seen³¹. Polymyalgia rheumatica is another condition that must be remembered in FM differential diagnosis, despite the typical involvement of the pelvic and scapular girdles and the significant elevation of erythrocyte sedimentation rate³². The fact that patients with polymyalgia rheumatica have a satisfactory and rapid response to low-dose oral corticosteroids can contribute to diagnostic elucidation. However, muscular ache, the predominant characteristic in this illness, might confuse a less attentive practitioner, particularly when muscular ache is diffuse.

In some cases, polymyositis (as well as other myositis) may be confused with FM. In this disease, it is important to remember that the main complaint is weakness rather than muscular pain. This weakness is characteristically insidious, symmetrical, and progressive, particularly at the scapular and pelvic girdles. It is painful in approximately 15% to 30% of cases. Patients may have arthralgia, similar to FM patients. Other clinical manifestations of muscular inflammatory disease, along with laboratory abnormalities, particularly high muscle enzymes, lead to the correct diagnosis³³.

Some autoimmune collagen diseases, particularly rheumatoid arthritis and systemic lupus erythematosus, may initially manifest as diffuse pain and fatigue, leading to FM misdiagnosis. In other situations, patients with established rheumatoid arthritis or systemic lupus erythematosus, with the various clinical and laboratory manifestations of these illnesses, may simultaneously present with a FM clinical picture. Despite the appropriate therapeutic response of the inflammation, painful complaints may remain. In these cases, synovitis disappears and inflammatory activity tests are normalized, with the concomitant maintenance of a FM clinical picture, coursing independently of the underlying disease. In one study, fibromyalgia coexistence was found in 12% of patients with rheumatoid arthritis³⁴. Not infrequently, Sjögren syndrome may have, as an initial manifestation, symptoms of diffuse and unspecific muscle pain accompanied by tiredness, making the diagnosis more difficult until laboratory parameters and other clinical characteristics are established³⁵.

Certain adverse reactions to drugs may cause diffuse myalgia, thus confusing clinical follow-up. Among these drugs, H₂ receptor blocking agents (used in peptic disease), fibrates, and statins (used in the treatment of dyslipidemia)³⁶ should be mentioned. Illicit drug users may experience similar reactions to a FM clinical picture,

particularly patients using cocaine and *Cannabis*. Patients with a history of alcoholism may also have muscle pain during either the alcohol abuse period or the withdrawal period³⁷.

Certain infections, in particular hepatitis C, human immunodeficiency virus (HIV), and Lyme disease, may cause diffuse muscular pain, making diagnosis more difficult, mainly when fever and other signs are not found³⁸. Serum analysis can contribute to elucidate some cases.

Patients who underwent long-term corticosteroid therapy, irrespective of the reason, can experience corticosteroid withdrawal, particularly if the drug cessation has been abrupt or inappropriate. These patients can experience diffuse muscle pain, rapidly responding to corticosteroid reintroduction³⁹.

Another condition that must be accounted for in FM differential diagnosis is paraneoplastic syndrome. Obviously, there are other signs directing the correct clinical reasoning in this picture. However, in rare situations, a paraneoplasm may manifest with a clinical picture similar to FM and issue an important diagnostic challenge, particularly in the case of bronchogenic carcinoma⁴⁰.

Chronic fatigue syndrome should also be included into the differential diagnosis. Low fever or recurring pharyngitis history, in addition to lymphadenopathy, can contribute to clarify the diagnosis. In chronic fatigue syndrome, tiredness predominates, while diffuse pain predominates in fibromyalgia⁴¹.

Possibly the greatest diagnostic challenge is differentiating FM from psychogenic rheumatism. In certain situations, the patient's clinical picture purely expresses a psychiatric disorder, particularly depression. Depression is known to occur more commonly in fibromyalgia than in controls or even in patients with rheumatoid arthritis⁴². About half of patients with FM have had a depression episode at some time in their lives. Many depression symptoms, such as tiredness, energy loss, discouragement, and sleep disturbances, are identical to those in FM. Maybe this is why many authors consider fibromyalgia as a depression manifestation. However, it is important to keep in mind that a large number of patients with fibromyalgia have neither depression nor any other psychiatric disorder component⁴³.

It is also important to note that sometimes patients with fibromyalgia have been misdiagnosed as having RSI/WRMD (repetition strain injury/work-related musculoskeletal disorders). In an investigation conducted in the Discipline of Rheumatology at Escola Paulista de Medicina, 103 patients with an initial diagnosis of repetition strain injury (RSI) or work-related musculoskeletal disorder (WRMD), all of them involved in a labor dispute, were studied. Among these 103 patients, 73

were found to meet diagnostic criteria for FM and they had no evidence of tissue injury⁴⁴.

Finally, RSI and WRMD cannot be widely pointed out as a disease, but they are acronyms embodying a heterogeneous group of musculoskeletal disorders having a work agent as one of the required causes for their appearance. It is important to point out that FM is not a RSI or a WRMD.

TREATMENT

To this day, the lack of knowledge about FM etiopathogenesis prevents a totally effective treatment.

FM treatment should be multidisciplinary, individualized, count on the patient's active participation, based on combined non-pharmacological and pharmacological therapy and should be designed according to the intensity and characteristics of symptoms. It is important that biopsychosocial issues involved in the context of illness are also considered.

As an initial part of treatment, patients are provided with basic information about FM and treatment options, and are instructed about pain control and self-control programs.

FM drug therapy, in addition to pain control, aims to induce a better quality sleep and treat associated symptoms, such as depression and anxiety.

Nonsteroidal anti-inflammatory drugs (NSAIDs) should not be used as first-line drugs in patients with FM⁴⁵. Although there is no evidence of inflammation, NSAIDs used in treating more prominent painful complaints act satisfactorily on FM-associated symptoms, such as headache and joint pain⁴⁶.

Corticosteroids are not a part of the therapeutic resources used in FM⁴⁷.

In Brazil, two out of three drugs approved by the Food and Drug Administration (FDA) to treat FM are available: pregabalin and duloxetine. Pregabalin is a calcium channel modulating agent, reducing the release of excitatory pain neurotransmitter release at nervous endings, particularly substance P and glutamate. Studies show significant relief of pain, fatigue, anxiety, and sleep disturbances by using this drug⁴⁸. Duloxetine is a serotonin and noradrenalin reuptake inhibitor that also has shown to be effective in reducing pain and improving functional capacity in FM regardless of the presence of depression⁴⁹.

Tricyclic antidepressants, especially amitriptylin and cyclobenzaprine, ingested two to three hours before bedtime, can be effective in decreasing the pain and improving the sleep, as well as enhancing the functional capacity^{50,51}.

Many other drugs were studied, but the results were usually less satisfactory⁵².

Non-pharmacological treatment has a crucial role in managing FM symptoms⁵³.

FM patients benefit from physical activity. Several reasons justify the encouragement of physical activity in this syndrome: increased levels of serotonin and other inhibitory neurotransmitters; increased GH (growth hormone) production; hypothalamic-pituitary-adrenal axis and autonomous nervous system regulation; increased capillary density; increased myoglobin count; and increased mitochondrial activity. All these changes contribute to improve pain, quality of sleep, fatigue, anxiety, and other symptoms. The possibility of socialization, depending on the circumstances, and the positive influence on some psychological aspects should also be taken into account. Physical activity has represented the most frequently employed and studied non-drug intervention for FM. However, the most appropriate type has not been established, as well as the optimal frequency and intensity⁴⁸.

Psychological aspects should always be evaluated and occasional disorders should be identified, aiming at real support and specialized help as necessary. Sleep disturbances should also be approached and treated⁵⁴.

In a recent literature review, involving mainly systematic reviews and meta-analysis publications over last 12 years, a lack of uniformity on the type of rehabilitation and exercise that should be employed in this chronic pain syndrome was verified. The forms of aerobic activity studied (walking, marching, cycling, rowing, etc.) were varied, as well as other types of exercising (stretching, isometric and isokinetic exercises, among other exercises for muscular strengthening), and various forms of hydrotherapy (aquatic respiratory exercises, deep water running, hydrogymnastics, swimming, among others). A great diversity in rehabilitation methods, such as coping, biofeedback, family education, cognitive-behavioral therapy, and manipulation and relaxation techniques was also found. Overall, the studies concluded that the effects were beneficial both from exercises and from different types of rehabilitation, even if involving heterogeneous modalities⁴⁸.

OCCUPATIONAL ASPECTS

Scientific literature offers no evidence that FM is occupational in origin. In the 1990's, some scientific papers related to occupational trauma and litigation involving FM were published⁵⁵⁻⁵⁷.

However, literature data are not enough to establish causality. The connection with repetition microtrauma has been suggested by some authors; however, these studies were either conducted with small samples of patients or documented with isolated case reports^{58,59}.

FM does not usually lead to incapacity to work. In cases of significant pain or fatigue, taking a few sick days can be considered. Nevertheless, litigious repetitive requests of temporary leaves have been observed in judicial practice, resulting in a long and continued absence of the patient. There is no systematic investigation of these lawsuits aiming to quantify how often this has occurred or how its economic dimension has impacted the social security system in Brazil. Data are empirical and observational. It cannot be denied that a protracted leave and the search for financial compensation in a chronic pain setting may be harmful, as investigations have demonstrated^{60,61}.

Of note, individuals with chronic pain are prone to persistence of painful symptomatology when they are involved in litigation⁶². In addition, there is increased malingering in a permissive system of judicial compensation^{63,64}.

It is debatable whether patients with fibromyalgia experience their symptoms worsened when their functions are performed under strict productivity criteria and pressured by time, such as in assembly lines. To date, the biomechanical limits of these patients have not been objectively defined, making it impossible to establish reduction, modification, or elimination criteria for any specific work setting. A worker with FM might keep performing his/her work, more slowly, with his/her productivity criteria or demands reconsidered.

Systematic reviews, such as that by Boocock et al., also establish that there is no current specific modification in work or labor intervention strategy that can be considered effective to solve the occupational issues involving these patients. The authors also mention the lack of research for several labor activities⁶⁵.

In FM, as in any other chronic pain syndrome, the sense of incapacity can be influenced by social and psychological factors. Work incapacity allegations may be vicious when psychological factors are predominant, or when pain or fatigue seems less severe than reported. The problem can become more pronounced when there is an interest in secondary gain. In settings where ensuring the absence of such interest is difficult to prove, the only solution is to keep a prolonged longitudinal follow-up, which improves doctor-patient relationship.

It is important to note that courts in the United States usually do not grant benefits to patients with FM⁶⁶.

These patients are prone to have their symptoms worsened when they present with associated psychological disorders. A study conducted in Brazil found that 30% of FM patients exhibited severe depression and 34%, moderate depression; this same study found that 70% of FM patients had traits of significant anxiety and 88% exhibited a high anxiety status⁶⁷.

Concerning depression, Brazilian labor laws have specific terms for this illness. However, it is imperative not to confuse depression with FM. A physician should not attempt to diagnose depression without having enough experience and without well-established diagnostic criteria.

If there is a significant level of depression, the possibility of a medical leave should be considered. However, the patient must not feel comfortable in this situation, due to the consequent creation of a vicious circle and no adherence to treatment.

Another noteworthy aspect is the possibility of malingering. In a controlled study involving FM patients, healthy controls and patients motivated to malingering, the American College of Rheumatology (ACR) criteria were found to be 80% accurate, with a good agreement and reproducibility level towards tender points and control points so as to distinguish malingerers. This study's results demonstrate that there is no malingering test and it is likely a number of malingerers and true FM patients could remain unidentified⁶⁸.

FINAL CONSIDERATIONS

FM is a syndrome with a prominently clinical diagnosis, characterized by chronic musculoskeletal pain associated with various symptoms. It may be confused with several rheumatic and non-rheumatic diseases. An individualized and multidisciplinary approach, combining drug and nondrug treatment, is required.

There is no scientific evidence determining that FM is caused by work. However, reasonable labor adaptations should be sought, including reduced time for task execution, among others, although labor modifications for this disease are not yet defined.

If the patient with fibromyalgia presents significant levels of pain or fatigue and does not respond to appropriate treatment, a short medical leave could be considered, although repetitive temporary leaves obtained by litigious means have been empirically observed, but they have no scientific support for their real requirement and their consequent economic repercussions. There is negative evidence for patients regarding litigation and long leaves. Prolonged leaves are justified in the case of associated severe depression.

When approaching patients with fibromyalgia, it is important to explain this chronic pain syndrome and the barriers against clinical improvement.

REFERENCES

1. Smythe HA, Moldofsky H. Two contributions to understanding of the "fibrositis" syndrome. *Bull Rheum Dis*. 1977;28:928-31.
2. Yunus MB, Masi AT, Calabro JJ. Primary fibromyalgia (fibrositis). Clinical study of 50 patients with matched normal controls. *Semin Arthritis Rheum*. 1981;11:151-71.

3. Valentine M. Aetiology of fibrositis: a review. *Ann Rheum Dis.* 1947;6:241-50.
4. Wolfe F, Smithe HA, Yunus MB, Bennett RM, Bombardier C, Goldenberg DL, et al. The American College of Rheumatology 1990 criteria for the classification of fibromyalgia. Report of the multicenter Criteria Committee. *Arthritis Rheum.* 1990;33:160-72.
5. Wolfe F, Clauw DJ, Fitzcharles MA, Goldenberg DL, Katz RS, Mease P, et al. The American College of Rheumatology preliminary diagnostic criteria for fibromyalgia and measurement of symptom severity. *Arthritis Care Res.* 2010;62:600-10.
6. Wolfe F, Ross K, Anderson J, Russell IJ, Herbert I. The prevalence and characteristics of fibromyalgia in the general population. *Arthritis Rheum.* 1995;38:19-28.
7. Forseth KO, Gran JT. The prevalence of fibromyalgia among women aged 20-49 years in Aredal, Norway. *Scand J Rheumatol.* 1992;21:74-8.
8. Senna ER, De Barros AL, Silva EO, Costa IF, Pereira LV, Ciconelli RM, et al. Prevalence of rheumatic diseases in Brazil: a study using the COPCORD approach. *J Rheumatol.* 2004;31:594-7.
9. Forseth KO, Gran JT, Husby GA. A population study of the incidence of fibromyalgia among women aged 26-55 years. *Br J Rheumatol.* 1997;36:1318-23.
10. Wolfe F. The relation between tender points and fibromyalgia symptom variables: evidence that fibromyalgia is not a discrete disorder in the clinic. *Ann Rheum Dis.* 1997;56: 268-71.
11. Berger A, Dukes E, Martin S, Edelsberg J, Oster G. Characteristics and healthcare costs of patients with fibromyalgia syndrome. *Int J Clin Pract.* 2007;61:1498-508.
12. Boonen A, Van Den Heuvel R, Van Tubergen A, Goossens M, Severens J, Van Der Heijde D, et al. Large differences in cost of illness and wellbeing between patients with fibromyalgia, chronic low back pain, or ankylosing spondylitis. *Ann Rheum Dis.* 2005;64:396-402.
13. Spaeth M. Epidemiology, costs, and the economic burden of fibromyalgia. *Arthritis Res Ther.* 2009;11:117-21.
14. Bradley LA, McKendree-Smith NL. Central nervous system mechanisms of pain in fibromyalgia and other musculoskeletal disorders: behavioral and psychologic treatment approaches. *Curr Opin Rheumatol.* 2002;14:45-51.
15. Buskila D, Neumann L. Fibromyalgia syndrome (FM) and nonarticular tenderness in relatives of patients with FM. *J Rheumatol.* 1997;24:941-4.
16. Arnold LM, Hudson JI, Hess EV, Ware AE, Fritz DA, Auchenbach MB, et al. Family study of fibromyalgia. *Arthritis Rheum.* 2004;50:944-52.
17. Crofford LJ, Demitrack MA. Evidence that abnormalities of central neurohormonal systems are key to understanding fibromyalgia and chronic fatigue syndrome. *Rheum Dis Clin North Am.* 1996;22:267-84.
18. Helfenstein M, Feldman D. Síndrome da fibromialgia: características clínicas e associações com outras síndromes disfuncionais. *Rev Bras Reumatol.* 2002;42:8-14.
19. Croft P, Schollum J, Silman A. Population study of tender point counts and pain as evidence of fibromyalgia. *BMJ.* 1994;309:696-9.
20. Cote KA, Moldofsky H. Sleep, daytime symptoms, and cognitive performance in patients with fibromyalgia. *J Rheumatol.* 1997;24:2014-23.
21. Zammit GK, Weiner J, Damato N, Sillup GP, McMillan CA. Quality of life in people with insomnia. *Sleep.* 1999;22:379-85.
22. Weidebach WFS. Fibromialgia: evidências de um substrato neurofisiológico. *Rev Assoc Med Bras.* 2002;48:291-5.
23. Raphael KG, Janal MN, Nayak S, Schwartz JE, Gallagher RM. Familial aggregation of depression in fibromyalgia: a community-based test of alternate hypotheses. *Pain.* 2004;110:449-60.
24. Nordahl HM, Stiles TC. Personality styles in patients with fibromyalgia, major depression and healthy controls. *Ann Gen Psychiatry.* 2007;6:9-13.
25. Da Costa PL, Dobkin M, Fitzcharles MA. The relationship between exercise participation and depressed mood in women with fibromyalgia. *Psychol Health Med.* 2001;6:301-11.
26. Berber JSS, Kupek E, Berber SC. Prevalência de depressão e sua relação com a qualidade de vida em pacientes com síndrome da fibromialgia. *Rev Bras Reumatol.* 2005;45:47-54.
27. Collado A, Alijotas B, Benitoc P, Alegred C, Romerae M, Sañudof I, et al. Documento de consenso sobre el diagnóstico y tratamiento de la fibromialgia en Cataluña. *Med Clin (Barc).* 2002;118:745-9.
28. Bazzichi L, Rossi A, Giuliano T, Feo F, Giacomelli C, Consensi A, et al. Association between thyroid autoimmunity and fibromyalgic disease severity. *Clin Rheumatol.* 2007;26:2115-20.
29. Sulea AA, Leow MKS. Association of polycystic ovary syndrome and fibromyalgia in a patient with primary hyperparathyroidism: a novel triad? *Clin Chem.* 2006;52:1208-9.
30. Erkal M Z, Wilde J, Bilgin Y, Akinci A, Demir E, Bödeker RH, et al. High prevalence of vitamin D deficiency, secondary hyperparathyroidism and generalized bone pain in Turkish immigrants in Germany: identification of risk factors. *Osteop Int.* 2006;17:1133-40.
31. Al-Allaf AW, Mole PA, Paterson CR and Pullar T. Bone health in patients with fibromyalgia. *Rheumatology.* 2003;42:1202-6.
32. Nothnagl T, Leeb B. Diagnosis, differential diagnosis and treatment of poly-myalgia rheumatica. *Drugs Aging.* 2006;23:391-402.
33. Dadabhoy D, Clauw DJ. Musculoskeletal signs and symptoms. The fibromyalgia syndrome. In: Klippel JH, editor. *Primer of the rheumatic diseases.* 13th ed. Philadelphia: Verlag; 2008. p. 87-93.
34. Martinez-Lavin M. Overlap of fibromyalgia with other medical conditions. *Curr Pain Headache Reports.* 2007;5:347-50.
35. Kassin SS, Moutsopoulos HM. Clinical manifestations and early diagnosis of Sjögren syndrome. *Arch Intern Med.* 2004;164:1275-84.
36. Mascitelli L, Pezzetta F, Goldstein MR. Detrimental effect of statin therapy in women with fibromyalgia. *Arch Intern Med.* 2008;168:1228-9.
37. Eisinger J. Alcohol, thiamin and fibromyalgia. *J Am Coll Nutr.* 1998;17:300-2.
38. Kozanoglu E, Canatarglu A, Abayli B, Colakoglu S, Goncu K. Fibromyalgia syndrome in patients with hepatitis C infection. *Rheumatol Int.* 2004;23:248-51.
39. Adler GK, Manfredsdottir VF, Rackow RM. Hypothalamic-pituitary-adrenal axis function in fibromyalgia and chronic fatigue syndrome. *Endocrinologist.* 2002;12:513-24.
40. Gopaluni S, Thomas LM, Shah C. When myopathy strikes in a cancer patient. *Gastrointest Cancer Res.* 2009;3:212-4.
41. Friedberg F, Jason LA. Chronic fatigue syndrome and fibromyalgia: clinical assessment and treatment. *J Clin Psychol.* 2001;57:433-55.
42. McBeth J, Silman AJ. The role of psychiatric disorders in fibromyalgia. *Curr Rheumatol Rep.* 2001;3:157-64.
43. Winfield JB. Does pain in fibromyalgia reflect somatization? *Arthritis Rheum.* 2001;44:751-3.
44. Helfenstein M. Prevalência da síndrome da fibromialgia em pacientes diagnosticados como portadores de lesões por esforços repetitivos (LER) [tese]. São Paulo: Escola Paulista de Medicina; 1997.
45. Buckhardt CS, Goldenberg D, Crofford L, Gerwin R, Gowans S, Kugel P, et al. Guideline for the management of fibromyalgia syndrome pain in adults and children. APS Clinical Practice Guidelines Series, n. 4. Glenview: American Pain Society; 2005.
46. Leventhal LJ. Management of fibromyalgia. *Ann Intern Med.* 1999;131:850-8.
47. Carville SF, Arendt-Nielsen S, Bliddal H, Blotman F, Branco JC, Buskila D, et al. EULAR evidence-based recommendations for the management of fibromyalgia syndrome. *Ann Rheum Dis.* 2008;67:536-41.
48. Clark P, Gentile MJ, Helfenstein M, Jannaut MJ, Liendo V, Rios C, et al. Tratamiento farmacológico y no farmacológico de la fibromialgia - síntesis de la mejor evidencia. *Drugs Today.* 2011;47(Suppl A):1-28.
49. Arnold LM, Hudson JI, Wang F, Wohlreich MM, Prakash A, Kajdasz DK, et al. Comparisons of the efficacy and safety of duloxetine for the treatment of fibromyalgia in patients with versus without major depressive disorder. *Clin J Pain.* 2009;25:461-8.
50. Tofferi JK, Jackson JL, O'Malley PG. Treatment of fibromyalgia with cyclo-benzaprine: a meta-analysis. *Arthritis Care Res.* 2004;51:9-13.
51. Nishishinya B, Urrutia G, Walitt B, Rodriguez A, Bonfill X, Alegre C, et al. Amitriptyline in the treatment of fibromyalgia: a systematic review of its efficacy. *Rheumatology.* 2008;47:1741-6.
52. Heymann RE, Paiva ES, Helfenstein M, Pollak DF, Martinez JE, Provenza JR, et al. Consenso brasileiro do tratamento da fibromialgia. *Rev Bras Reumatol.* 2010;50:56-66.
53. Sim J, Adams N. Systematic review of randomized controlled trials of nonpharmacological interventions for fibromyalgia. *Clin J Pain.* 2002;18:324-36.
54. Roizenblatt S, Neto NS, Tufik S. Sleep disorders and fibromyalgia. *Curr Pain Headache Rep.* 2011;15:347-57.
55. Whorton D, Weisenberger BI, Milroy WC. Does fibromyalgia qualify as a work-related illness or injury? *J Occup Environ Med.* 1992;34:968-72.
56. Bruusgaard D, Evensen AR, Bjerkedal T. Fibromyalgia - a new cause for disability pension. *Scand J Soc Med.* 1993;21:116-93.
57. Bennett RM. Disabling fibromyalgia: appearance versus reality. *J Rheumatol.* 1993;11:1821-5.
58. Romano TJ. Clinical experiences with post-traumatic fibromyalgia syndrome. *West Virg Med J.* 1990;86:198-202.
59. Moldofsky H, Wong MTH, Lue FA. Litigation, sleep, symptoms and disabilities in postaccident pain (fibromyalgia). *J Rheumatol.* 1993;20:1935-40.
60. Rainville JR, Sobel JB, Hartigan C. The effect of compensation involvement on the reporting of pain and disability by patients referred for rehabilitation of chronic low back pain. *Spine.* 1997;22:2016-24.
61. Rohling ML, Binder LM, Langhinrichsen-Rohling JL. Money matters: a meta-analytic review of the association between financial compensation and the experience and treatment of chronic pain. *Health Psychol.* 1995;14:537-47.
62. Tait RC, Chibnall JT, Richardson WD. Litigation and employment status: effects on patients with chronic pain. *Pain.* 1990;43:37-46.
63. Figlio K. How does illness mediate social relations? Workmen's compensation and medico-legal practices, 1890-1940. In: Wright P, Treacher A, editors. *The problem of medical knowledge.* 2nd ed. Edinburgh: Edinburgh University Press; 1982. p.174-224.
64. Bell DS. "Repetition strain injury": an iatrogenic epidemic of simulated injury. *Med J Aust.* 1989;151:280-4.

65. Boocock MG, McNair PJ, Larmer PJ, Armstrong B, Collier J, Simmonds M, et al. Interventions for the prevention and management of neck/upper extremity musculoskeletal conditions: a systematic review. *Occup Environ Med.* 2007;64:291-303.
66. Wolfe F, Potter J. Fibromyalgia and work disability: is fibromyalgia a disabling disorder? *Rheum Dis Clin North Am.* 1996;22:369-91.
67. Helfenstein M, Feldman D. Prevalência da síndrome da fibromialgia em pacientes diagnosticados como portadores de lesões por esforços repetitivos (LER). *Rev Bras Reumatol.* 1998;38:71-7.
68. Khostanteen I, Tunks ER, Goldsmith CH, Ennis J. Fibromyalgia: can one distinguish it from simulation? An observer-blind controlled study. *J Rheumatol.* 2000;27:2671-6.